

TITLE**PLANT MYB TRANSCRIPTION FACTOR HOMOLOGS**

This application claims the benefit of U.S. Provisional Application No. 60/110,609, filed December 2, 1998.

5

FIELD OF THE INVENTION

This invention is in the field of plant molecular biology. More specifically, this invention pertains to nucleic acid fragments encoding Myb-related transcription factors in plants and seeds.

BACKGROUND OF THE INVENTION

10

Improvement of crop plants for a variety of traits, including disease and pest resistance, and grain quality improvements such as oil, starch or protein composition, can be achieved by introducing new or modified genes (transgenes) into the plant genome.

Transcriptional activation of genes, including transgenes, is in general controlled by the promoter through a complex set of protein/DNA and protein/protein interactions. Promoters can impart patterns of expression that are either constitutive or limited to specific tissues or times during development.

15

Transcriptional activation is primarily mediated through transcription factors that interact with enhancer and promoter elements. Binding of transcription factors to such DNA elements constitutes a crucial step in transcriptional initiation. Each transcription factor binds to its specific binding sequence in a promoter and activates expression of the linked coding region through interactions with coactivators and/or proteins that are a part of the transcription complex.

20

Several plant genes have been identified that appear to encode transcription factors structurally related to the cMyb protooncogene family of mammals. Central to the similarities shared by these proteins is the Myb repeat DNA-binding domain containing conserved tryptophan residues at certain positions, and a helix-turn-helix-related domain. Generally, Myb-related proteins from plants contain two of these repeats, R2 and R3 (Kranz et al. (1998) *Plant J* 16:263-276), though proteins having only one repeat have been identified (e.g., Feldbrugge et al. (1997) *Plant J* 11:1079-1093). These Myb-related genes appear to encode a large family of plant transcription factors that are involved in a diversity of gene regulation. For example, plant Myb-related genes have been shown to regulate anthocyanin biosynthesis in maize and phenylpropanoid metabolism, disease resistance (WO9813486-A1), expression of gibberellin-regulated genes (WO9700961-A1), expression of stress-related genes (WO9916878-A1), active carbohydrate secretion and flavonol metabolism in antirrhinum flowers (Jackson et al. (1992) *Plant Cell* 3(2):115-125). The first plant transcription activator gene described at the molecular level was the maize *c1* gene which encodes a Myb protein (Paz-Ares et al. (1987) *EMBO J* 16:3553-3558) involved in

25

30

35

regulating anthocyanin biosynthesis by trans-activating genes such as *c2*, *A1* and *Bz1* which encode enzymes involved in the pathway.

There is a great deal of interest in identifying the genes that encode proteins involved in transcriptional regulation in plants. These genes may be used in plant cells to control gene expression constitutively, in specific tissues or at various times during development. Accordingly, the availability of nucleic acid sequences encoding all or a portion of a Myb-related transcription factor would facilitate studies to better understand gene regulation in plants and provide genetic tools to enhance or otherwise alter the expression of genes controlled by Myb-related transcription factors.

SUMMARY OF THE INVENTION

The present invention relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 50 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a corn Myb-related transcription factor polypeptide of SEQ ID NO:2, a rice Myb-related transcription factor polypeptide of SEQ ID NO:12, and a wheat Myb-related transcription factor polypeptide of SEQ ID NO:56. The present invention also relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 50 amino acids that has at least 85% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a corn Myb-related transcription factor polypeptide of SEQ ID NO:8, and a soybean Myb-related transcription factor polypeptide of SEQ ID NO:28. The present invention also relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 50 amino acids that has at least 90% identity based on the Clustal method of alignment when compared to a rice Myb-related transcription factor polypeptide of SEQ ID NO:16. The present invention also relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 50 amino acids that has at least 95% identity based on the Clustal method of alignment when compared to a soybean Myb-related transcription factor polypeptide of SEQ ID NO:52. The present invention also relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 100 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a corn Myb-related transcription factor polypeptide of SEQ ID NO:6, a rice Myb-related transcription factor polypeptide of SEQ ID NO:14, a soybean Myb-related transcription factor polypeptide of SEQ ID NO:50, and a wheat Myb-related transcription factor polypeptide of SEQ ID NO:58. The present invention also relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 100 amino acids that has at least 85% identity based on the Clustal method of alignment when compared to a wheat Myb-related transcription factor polypeptide of SEQ ID NO:60. The present invention also relates to

isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 100 amino acids that has at least 90% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a corn Myb-related transcription factor polypeptide of SEQ ID NO:4, a corn Myb-related transcription factor polypeptide of SEQ ID NO:10, a rice Myb-related transcription factor polypeptide of SEQ ID NO:22, a rice Myb-related transcription factor polypeptide of SEQ ID NO:24, and a wheat Myb-related transcription factor polypeptide of SEQ ID NO:62. The present invention also relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 100 amino acids that has at least 95% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a rice Myb-related transcription factor polypeptide of SEQ ID NO:18 and a rice Myb-related transcription factor polypeptide of SEQ ID NO:20. The present invention also relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 150 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a rice Myb-related transcription factor polypeptide of SEQ ID NO:26, a soybean Myb-related transcription factor polypeptide of SEQ ID NO:34, a soybean Myb-related transcription factor polypeptide of SEQ ID NO:38, a soybean Myb-related transcription factor polypeptide of SEQ ID NO:40, a soybean Myb-related transcription factor polypeptide of SEQ ID NO:42, a soybean Myb-related transcription factor polypeptide of SEQ ID NO:48, and a soybean Myb-related transcription factor polypeptide of SEQ ID NO:54. The present invention also relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 150 amino acids that has at least 85% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of a soybean Myb-related transcription factor polypeptide of SEQ ID NO:32, a soybean Myb-related transcription factor polypeptide of SEQ ID NO:44, and a soybean Myb-related transcription factor polypeptide of SEQ ID NO:46. The present invention also relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 200 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a soybean Myb-related transcription factor polypeptide of SEQ ID NO:36. The present invention also relates to isolated polynucleotides comprising a nucleotide sequence encoding a first polypeptide of at least 200 amino acids that has at least 85% identity based on the Clustal method of alignment when compared to a soybean Myb-related transcription factor polypeptide of SEQ ID NO:30. The present invention also relates to an isolated polynucleotide comprising the complement of the nucleotide sequences described above.

It is preferred that the isolated polynucleotides of the claimed invention consist of a nucleic acid sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11,

13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61 that codes for the polypeptide selected from the group consisting of SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, and 62. The present invention also relates to an isolated polynucleotide comprising a nucleotide sequences of at least one of 60 (preferably at least one of 40, most preferably at one least of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61 and the complement of such nucleotide sequences.

The present invention relates to a chimeric gene comprising an isolated polynucleotide of the present invention operably linked to suitable regulatory sequences.

The present invention relates to an isolated host cell comprising a chimeric gene of the present invention or an isolated polynucleotide of the present invention. The host cell may be eukaryotic, such as a yeast or a plant cell, or prokaryotic, such as a bacterial cell. The present invention also relates to a virus, preferably a baculovirus, comprising an isolated polynucleotide of the present invention or a chimeric gene of the present invention.

The present invention relates to a process for producing an isolated host cell comprising a chimeric gene of the present invention or an isolated polynucleotide of the present invention, the process comprising either transforming or transfecting an isolated compatible host cell with a chimeric gene or isolated polynucleotide of the present invention.

The present invention relates to a Myb-related transcription factor polypeptide of at least 50 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:2, 12, and 56. The present invention also relates to a Myb-related transcription factor polypeptide of at least 50 amino acids that has at least 85% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:8 and 28. The present invention also relates to a Myb-related transcription factor polypeptide of at least 50 amino acids that has at least 90% identity based on the Clustal method of alignment when compared to a rice Myb-related transcription factor polypeptide of SEQ ID NO:16. The present invention also relates to a Myb-related transcription factor polypeptide of at least 50 amino acids that has at least 95% identity based on the Clustal method of alignment when compared to a soybean Myb-related transcription factor polypeptide of SEQ ID NO:52. The present invention also relates to a Myb-related transcription factor polypeptide of at least 100 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:6, 14, 50, and 58. The present invention also relates to a Myb-related transcription factor polypeptide of at least 100 amino acids that has at least 85% identity based on the Clustal method of alignment when compared to a wheat Myb-related

transcription factor polypeptide of SEQ ID NO:60. The present invention also relates a Myb-related transcription factor polypeptide of at least 100 amino acids that has at least 90% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:4, 10, 22, 24, and 62. The present invention also relates to a Myb-related transcription factor polypeptide of at least 100 amino acids that has at least 95% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:18 and 20. The present invention also relates to a Myb-related transcription factor polypeptide of at least 150 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:26, 34, 38, 40, 42, 48, and 54. The present invention also relates to a Myb-related transcription factor polypeptide of at least 150 amino acids that has at least 85% identity based on the Clustal method of alignment when compared to a polypeptide selected from the group consisting of SEQ ID NOs:32, 44, and 46. The present invention also relates to a Myb-related transcription factor polypeptide of at least 200 amino acids that has at least 80% identity based on the Clustal method of alignment when compared to a soybean Myb-related transcription factor polypeptide of SEQ ID NO:36. The present invention also relates to a Myb-related transcription factor polypeptide of at least 200 amino acids that has at least 85% identity based on the Clustal method of alignment when compared to a soybean Myb-related transcription factor polypeptide of SEQ ID NO:30.

The present invention relates to a method of selecting an isolated polynucleotide that affects the level of expression of a Myb-related transcription factor polypeptide in a host cell, preferably a plant cell, the method comprising the steps of:

- constructing an isolated polynucleotide of the present invention or an isolated chimeric gene of the present invention;
- introducing the isolated polynucleotide or the isolated chimeric gene into a host cell;
- measuring the level of a Myb-related transcription factor polypeptide in the host cell containing the isolated polynucleotide; and
- comparing the level of a Myb-related transcription factor polypeptide in the host cell containing the isolated polynucleotide with the level of a Myb-related transcription factor polypeptide in a host cell that does not contain the isolated polynucleotide.

The present invention relates to a method of obtaining a nucleic acid fragment encoding a substantial portion of a Myb-related transcription factor polypeptide gene, preferably a plant Myb-related transcription factor polypeptide gene, comprising the steps of: synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3,

5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61 and the complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the oligonucleotide primer. The amplified nucleic acid fragment preferably will encode all or a portion of a Myb-related transcription factor amino acid sequence.

The present invention also relates to a method of obtaining a nucleic acid fragment encoding all or a substantial portion of the amino acid sequence encoding a Myb-related transcription factor polypeptide comprising the steps of: probing a cDNA or genomic library with an isolated polynucleotide of the present invention; identifying a DNA clone that hybridizes with an isolated polynucleotide of the present invention; isolating the identified DNA clone; and sequencing the cDNA or genomic fragment that comprises the isolated DNA clone.

BRIEF DESCRIPTION OF THE DRAWING AND SEQUENCE DESCRIPTIONS

The invention can be more fully understood from the following detailed description and the accompanying drawing and Sequence Listing which form a part of this application.

Figure 1 depicts the amino acid alignment between the Myb-related transcription factor encoded by the nucleotide sequences derived from corn clone cta1n.pk0079.e9 (SEQ ID NO:10), contig assembled from rice clones rr1.pk0027.g9 and rr1.pk077.n9 (SEQ ID NO:14), rice clone rl0n.pk082.c13 (SEQ ID NO:26), soybean clone sfl1.pk0032.g4 (SEQ ID NO:30), soybean clone sfl1.pk0086.a9 (SEQ ID NO:32), soybean clone sfl1.pk0091.a2 (SEQ ID NO:34), soybean clone sfl1.pk0091.a2 (SEQ ID NO:36), soybean clone sfl1.pk0003.a3 (SEQ ID NO:42), soybean clone srr3c.pk002.k6 (SEQ ID NO:44), soybean clone ses9c.pk002.o16 (SEQ ID NO:46), soybean clone sl2.pk127.e14 (SEQ ID NO:48), soybean clone src3c.pk010.i22 (SEQ ID NO:50), soybean clone sgs4c.pk004.j24 (SEQ ID NO:54), and a Myb-related transcription factor-encoding nucleic acid fragment from *Pisum sativum* (NCBI General Identification No. 1841475) (SEQ ID NO:63). Amino acids which are conserved among all and at least two sequences with an amino acid at that position are indicated with an asterisk (*) above them. Dashes are used by the program to maximize alignment of the sequences.

Table 1 lists the polypeptides that are described herein, the designation of the cDNA clones that comprise the nucleic acid fragments encoding polypeptides representing all or a substantial portion of these polypeptides, and the corresponding identifier (SEQ ID NO:) as used in the attached Sequence Listing. Table 1 also identifies the cDNA clones as individual ESTs ("EST"), the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS"), contigs assembled from two or more ESTs ("Contig"), contigs assembled from an FIS and one or more ESTs ("Contig*"), or sequences encoding the entire protein derived from an FIS, a contig, or an FIS and PCR ("CGS"). Nucleotide

SEQ ID NOs:7, 11, 23, 27, 51, and 55 correspond to nucleotide SEQ ID NOs:1, 3, 9, 5, 11, and 7, respectively, presented in U.S. Provisional Application No. 60/110,609, filed December 2, 1998. Amino acid SEQ ID NOs:8, 12, 24, 28, 52, and 56 correspond to amino acid SEQ ID NOs: 2, 4, 10, 6, 12, and 8, respectively presented in U.S. Provisional Application No. 60/110,609, filed December 2, 1998. The sequence descriptions and Sequence Listing attached hereto comply with the rules governing nucleotide and/or amino acid sequence disclosures in patent applications as set forth in 37 C.F.R. §1.821-1.825.

TABLE 1

Myb-related Transcription Factors

Protein	Clone Designation	Status	SEQ ID NO:	
			(Nucleotide)	(Amino Acid)
Myb-related Transcription Factor (Corn)	Contig of: csi1n.pk0008.b5 csi1n.pk0028.h7	Contig	1	2
Myb-related Transcription Factor (Corn)	Contig of: p0008.cb3ld06r p0026.ccrbd36rb	Contig	3	4
Myb-related Transcription Factor (Corn)	chpc8.pk0002.d9	EST	5	6
Myb-related Transcription Factor (Corn)	cta1n.pk0079.e9	EST	7	8
Myb-related Transcription Factor (Corn)	cta1n.pk0079.e9	CGS	9	10
Myb-related Transcription Factor (Rice)	rr1.pk077.n9	EST	11	12
Myb-related Transcription Factor (Rice)	Contig of: rr1.pk0027.g9 rr1.pk077.n9	CGS	13	14
Myb-related Transcription Factor (Rice)	rr1.pk088.p6	EST	15	16
Myb-related Transcription Factor (Rice)	rr1.pk0037.g7	EST	17	18
Myb-related Transcription Factor (Rice)	rds3c.pk002.c6	EST	19	20
Myb-related Transcription Factor (Rice)	Contig of: rlr24.pk0090.f5 rlr48.pk0012.c11	Contig	21	22
Myb-related Transcription Factor (Rice)	rl0n.pk082.c13	EST	23	24
Myb-related Transcription Factor (Rice)	rl0n.pk082.c13	CGS	25	26
Myb-related Transcription Factor (Soybean)	sfl1.pk0032.g4	EST	27	28

Protein	Clone Designation	Status	SEQ ID NO:	
			(Nucleotide)	(Amino Acid)
Myb-related Transcription Factor (Soybean)	sfl1.pk0032.g4	CGS	29	30
Myb-related Transcription Factor (Soybean)	sfl1.pk0086.a9	CGS	31	32
Myb-related Transcription Factor (Soybean)	sfl1.pk0091.a2	CGS	33	34
Myb-related Transcription Factor (Soybean)	sfl1.pk0105.e6	CGS	35	36
Myb-related Transcription Factor (Soybean)	sfl1.pk125.p19	FIS	37	38
Myb-related Transcription Factor (Soybean)	se6.pk0048.a12	FIS	39	40
Myb-related Transcription Factor (Soybean)	sfl1.pk0003.a3	CGS	41	42
Myb-related Transcription Factor (Soybean)	srr3c.pk002.k6	CGS	43	44
Myb-related Transcription Factor (Soybean)	ses9c.pk002.o16	CGS	45	46
Myb-related Transcription Factor (Soybean)	sl2.pk127.e14	CGS	47	48
Myb-related Transcription Factor (Soybean)	src3c.pk010.i22	CGS	49	50
Myb-related Transcription Factor (Soybean)	sgs4c.pk004.j24	EST	51	52
Myb-related Transcription Factor (Soybean)	sgs4c.pk004.j24	CGS	53	54
Myb-related Transcription Factor (Wheat)	wr1.pk0139.g11	EST	55	56
Myb-related Transcription Factor (Wheat)	wr1.pk0139.g11	FIS	57	58
Myb-related Transcription Factor (Wheat)	wdk3c.pk006.n12	EST	59	60
Myb-related Transcription Factor (Wheat)	wlm1.pk0027.a5	EST	61	62

The Sequence Listing contains the one letter code for nucleotide sequence characters and the three letter codes for amino acids as defined in conformity with the IUPAC-IUBMB standards described in *Nucleic Acids Res.* 13:3021-3030 (1985) and in the *Biochemical J.* 219 (No. 2):345-373 (1984) which are herein incorporated by reference. The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

DETAILED DESCRIPTION OF THE INVENTION

In the context of this disclosure, a number of terms shall be utilized. As used herein, a “polynucleotide” is a nucleotide sequence such as a nucleic acid fragment. A polynucleotide may be a polymer of RNA or DNA that is single- or double-stranded, that optionally
5 contains synthetic, non-natural or altered nucleotide bases. A polynucleotide in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA, synthetic DNA, or mixtures thereof. An isolated polynucleotide of the present invention may include at least one of 60 contiguous nucleotides, preferably at least one of 40
10 contiguous nucleotides, most preferably one of at least 30 contiguous nucleotides, of the nucleic acid sequence of the SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, 61, or the complement of such sequences.

As used herein, “contig” refers to a nucleotide sequence that is assembled from two or more constituent nucleotide sequences that share common or overlapping regions of
15 sequence homology. For example, the nucleotide sequences of two or more nucleic acid fragments can be compared and aligned in order to identify common or overlapping sequences. Where common or overlapping sequences exist between two or more nucleic acid fragments, the sequences (and thus their corresponding nucleic acid fragments) can be assembled into a single contiguous nucleotide sequence.

As used herein, “substantially similar” refers to nucleic acid fragments wherein changes in one or more nucleotide bases results in substitution of one or more amino acids, but do not affect the functional properties of the polypeptide encoded by the nucleotide
20 sequence. “Substantially similar” also refers to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate alteration of gene expression by gene silencing through for example antisense or co-suppression technology. “Substantially similar” also refers to modifications of the nucleic
25 acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially affect the functional properties of the resulting transcript vis-à-vis the ability to mediate gene silencing or alteration of the functional
30 properties of the resulting protein molecule. It is therefore understood that the invention encompasses more than the specific exemplary nucleotide or amino acid sequences and includes functional equivalents thereof.

Substantially similar nucleic acid fragments may be selected by screening nucleic acid fragments representing subfragments or modifications of the nucleic acid fragments of the
35 instant invention, wherein one or more nucleotides are substituted, deleted and/or inserted, for their ability to affect the level of the polypeptide encoded by the unmodified nucleic acid fragment in a plant or plant cell. For example, a substantially similar nucleic acid fragment representing at least one of 30 contiguous nucleotides derived from the instant nucleic acid

fragment can be constructed and introduced into a plant or plant cell. The level of the polypeptide encoded by the unmodified nucleic acid fragment present in a plant or plant cell exposed to the substantially similar nucleic fragment can then be compared to the level of the polypeptide in a plant or plant cell that is not exposed to the substantially similar nucleic acid fragment.

For example, it is well known in the art that antisense suppression and co-suppression of gene expression may be accomplished using nucleic acid fragments representing less than the entire coding region of a gene, and by nucleic acid fragments that do not share 100% sequence identity with the gene to be suppressed. Moreover, alterations in a nucleic acid fragment which result in the production of a chemically equivalent amino acid at a given site, but do not effect the functional properties of the encoded polypeptide, are well known in the art. Thus, a codon for the amino acid alanine, a hydrophobic amino acid, may be substituted by a codon encoding another less hydrophobic residue, such as glycine, or a more hydrophobic residue, such as valine, leucine, or isoleucine. Similarly, changes which result in substitution of one negatively charged residue for another, such as aspartic acid for glutamic acid, or one positively charged residue for another, such as lysine for arginine, can also be expected to produce a functionally equivalent product. Nucleotide changes which result in alteration of the N-terminal and C-terminal portions of the polypeptide molecule would also not be expected to alter the activity of the polypeptide. Each of the proposed modifications is well within the routine skill in the art, as is determination of retention of biological activity of the encoded products. Consequently, an isolated polynucleotide comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61 and the complement of such nucleotide sequences may be used in methods of selecting an isolated polynucleotide that affects the expression of a polypeptide in a plant cell. A method of selecting an isolated polynucleotide that affects the level of expression of a polypeptide (such as a Myb-related transcription factor) in a host cell (eukaryotic, such as plant or yeast, prokaryotic such as bacterial, or viral) may comprise the steps of: constructing an isolated polynucleotide of the present invention or an isolated chimeric gene of the present invention; introducing the isolated polynucleotide or the isolated chimeric gene into a host cell; measuring the level of a polypeptide in the host cell containing the isolated polynucleotide; and comparing the level of a polypeptide in the host cell containing the isolated polynucleotide with the level of a polypeptide in a host cell that does not contain the isolated polynucleotide.

Moreover, substantially similar nucleic acid fragments may also be characterized by their ability to hybridize. Estimates of such homology are provided by either DNA-DNA or

DNA-RNA hybridization under conditions of stringency as is well understood by those skilled in the art (Hames and Higgins, Eds. (1985) *Nucleic Acid Hybridisation*, IRL Press, Oxford, U.K.). Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly related organisms, to highly similar fragments, such as genes that duplicate functional enzymes from closely related organisms. Post-hybridization washes determine stringency conditions. One set of preferred conditions uses a series of washes starting with 6X SSC, 0.5% SDS at room temperature for 15 min, then repeated with 2X SSC, 0.5% SDS at 45°C for 30 min, and then repeated twice with 0.2X SSC, 0.5% SDS at 50°C for 30 min. A more preferred set of stringent conditions uses higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 min washes in 0.2X SSC, 0.5% SDS was increased to 60°C. Another preferred set of highly stringent conditions uses two final washes in 0.1X SSC, 0.1% SDS at 65°C.

Substantially similar nucleic acid fragments of the instant invention may also be characterized by the percent identity of the amino acid sequences that they encode to the amino acid sequences disclosed herein, as determined by algorithms commonly employed by those skilled in this art. Suitable nucleic acid fragments (isolated polynucleotides of the present invention) encode polypeptides that are at least about 70% identical, most preferably at least about 80% identical to the amino acid sequences reported herein. Preferred nucleic acid fragments encode amino acid sequences that are at least about 85% identical to the amino acid sequences reported herein. More preferred nucleic acid fragments encode amino acid sequences that are at least about 90% identical to the amino acid sequences reported herein. Most preferred are nucleic acid fragments that encode amino acid sequences that are at least about 95% identical to the amino acid sequences reported herein. Suitable nucleic acid fragments not only have the above homologies but typically encode a polypeptide having at least about 50 amino acids, preferably at least about 100 amino acids, more preferably at least about 150 amino acids, still more preferably at least about 200 amino acids, and most preferably at least about 250 amino acids. Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS*. 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5.

A "substantial portion" of an amino acid or nucleotide sequence comprises an amino acid or a nucleotide sequence that is sufficient to afford putative identification of the protein or gene that the amino acid or nucleotide sequence comprises. Amino acid and nucleotide sequences can be evaluated either manually by one skilled in the art, or by using computer-

based sequence comparison and identification tools that employ algorithms such as BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/). In general, a sequence of ten or more contiguous amino acids or thirty or more contiguous nucleotides is necessary in order to putatively identify a polypeptide or nucleic acid sequence as homologous to a known protein or gene. Moreover, with respect to nucleotide sequences, gene-specific oligonucleotide probes comprising 30 or more contiguous nucleotides may be used in sequence-dependent methods of gene identification (e.g., Southern hybridization) and isolation (e.g., *in situ* hybridization of bacterial colonies or bacteriophage plaques). In addition, short oligonucleotides of 12 or more nucleotides may be used as amplification primers in PCR in order to obtain a particular nucleic acid fragment comprising the primers. Accordingly, a "substantial portion" of a nucleotide sequence comprises a nucleotide sequence that will afford specific identification and/or isolation of a nucleic acid fragment comprising the sequence. The instant specification teaches amino acid and nucleotide sequences encoding polypeptides that comprise one or more particular plant proteins. The skilled artisan, having the benefit of the sequences as reported herein, may now use all or a substantial portion of the disclosed sequences for purposes known to those skilled in this art. Accordingly, the instant invention comprises the complete sequences as reported in the accompanying Sequence Listing, as well as substantial portions of those sequences as defined above.

"Codon degeneracy" refers to divergence in the genetic code permitting variation of the nucleotide sequence without effecting the amino acid sequence of an encoded polypeptide. Accordingly, the instant invention relates to any nucleic acid fragment comprising a nucleotide sequence that encodes all or a substantial portion of the amino acid sequences set forth herein. The skilled artisan is well aware of the "codon-bias" exhibited by a specific host cell in usage of nucleotide codons to specify a given amino acid. Therefore, when synthesizing a nucleic acid fragment for improved expression in a host cell, it is desirable to design the nucleic acid fragment such that its frequency of codon usage approaches the frequency of preferred codon usage of the host cell.

"Synthetic nucleic acid fragments" can be assembled from oligonucleotide building blocks that are chemically synthesized using procedures known to those skilled in the art. These building blocks are ligated and annealed to form larger nucleic acid fragments which may then be enzymatically assembled to construct the entire desired nucleic acid fragment. "Chemically synthesized", as related to nucleic acid fragment, means that the component nucleotides were assembled *in vitro*. Manual chemical synthesis of nucleic acid fragments may be accomplished using well established procedures, or automated chemical synthesis can be performed using one of a number of commercially available machines. Accordingly, the nucleic acid fragments can be tailored for optimal gene expression based on optimization of nucleotide sequence to reflect the codon bias of the host cell. The skilled artisan

appreciates the likelihood of successful gene expression if codon usage is biased towards those codons favored by the host. Determination of preferred codons can be based on a survey of genes derived from the host cell where sequence information is available.

“Gene” refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence. “Native gene” refers to a gene as found in nature with its own regulatory sequences. “Chimeric gene” refers any gene that is not a native gene, comprising regulatory and coding sequences that are not found together in nature.

Accordingly, a chimeric gene may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that found in nature.

“Endogenous gene” refers to a native gene in its natural location in the genome of an organism. A “foreign” gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric genes. A “transgene” is a gene that has been introduced into the genome by a transformation procedure.

“Coding sequence” refers to a nucleotide sequence that codes for a specific amino acid sequence. “Regulatory sequences” refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

“Promoter” refers to a nucleotide sequence capable of controlling the expression of a coding sequence or functional RNA. In general, a coding sequence is located 3' to a promoter sequence. The promoter sequence consists of proximal and more distal upstream elements, the latter elements often referred to as enhancers. Accordingly, an “enhancer” is a nucleotide sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic nucleotide segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters which cause a nucleic acid fragment to be expressed in most cell types at most times are commonly referred to as “constitutive promoters”. New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences

have not been completely defined, nucleic acid fragments of different lengths may have identical promoter activity.

The “translation leader sequence” refers to a nucleotide sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the primary transcript to mRNA, mRNA stability or translation efficiency. Examples of translation leader sequences have been described (Turner and Foster (1995) *Mol. Biotechnol.* 3:225-236).

The “3’ non-coding sequences” refer to nucleotide sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3’ end of the mRNA precursor. The use of different 3’ non-coding sequences is exemplified by Ingelbrecht et al. (1989) *Plant Cell* 1:671-680.

“RNA transcript” refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from posttranscriptional processing of the primary transcript and is referred to as the mature RNA. “Messenger RNA (mRNA)” refers to the RNA that is without introns and that can be translated into polypeptide by the cell. “cDNA” refers to a double-stranded DNA that is complementary to and derived from mRNA. “Sense” RNA refers to an RNA transcript that includes the mRNA and so can be translated into a polypeptide by the cell. “Antisense RNA” refers to an RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (see U.S. Patent No. 5,107,065, incorporated herein by reference). The complementarity of an antisense RNA may be with any part of the specific nucleotide sequence, i.e., at the 5’ non-coding sequence, 3’ non-coding sequence, introns, or the coding sequence. “Functional RNA” refers to sense RNA, antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes.

The term “operably linked” refers to the association of two or more nucleic acid fragments on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in sense or antisense orientation.

The term “expression”, as used herein, refers to the transcription and stable accumulation of sense (mRNA) or antisense RNA derived from the nucleic acid fragment of the invention. Expression may also refer to translation of mRNA into a polypeptide.

“Antisense inhibition” refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein. “Overexpression” refers to the production of a gene product in transgenic organisms that exceeds levels of production in normal or non-transformed organisms. “Underexpression” refers to the production of a gene product in transgenic organisms at levels below that of levels of production in normal or non-transformed organisms. “Co-suppression” refers to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar foreign or endogenous genes (U.S. Patent No. 5,231,020, incorporated herein by reference).

“Altered levels” refers to the production of gene product(s) in transgenic organisms in amounts or proportions that differ from that of normal or non-transformed organisms.

“Mature” protein refers to a post-translationally processed polypeptide; i.e., one from which any pre- or propeptides present in the primary translation product have been removed.

“Precursor” protein refers to the primary product of translation of mRNA; i.e., with pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

A “chloroplast transit peptide” is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the chloroplast or other plastid types present in the cell in which the protein is made. “Chloroplast transit sequence” refers to a nucleotide sequence that encodes a chloroplast transit peptide. A “signal peptide” is an amino acid sequence which is translated in conjunction with a protein and directs the protein to the secretory system (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53). If the protein is to be directed to a vacuole, a vacuolar targeting signal (*supra*) can further be added, or if to the endoplasmic reticulum, an endoplasmic reticulum retention signal (*supra*) may be added. If the protein is to be directed to the nucleus, any signal peptide present should be removed and instead a nuclear localization signal included (Raikhel (1992) *Plant Phys.* 100:1627-1632).

“Transformation” refers to the transfer of a nucleic acid fragment into the genome of a host organism, resulting in genetically stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as “transgenic” organisms. Examples of methods of plant transformation include *Agrobacterium*-mediated transformation (De Blaere et al. (1987) *Meth. Enzymol.* 143:277) and particle-accelerated or “gene gun” transformation technology (Klein et al. (1987) *Nature (London)* 327:70-73; U.S. Patent No. 4,945,050, incorporated herein by reference).

Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook et al. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989 (hereinafter “Maniatis”).

Nucleic acid fragments encoding at least a portion of several Myb-related transcription factors have been isolated and identified by comparison of random plant cDNA sequences to public databases containing nucleotide and protein sequences using the BLAST algorithms well known to those skilled in the art. The nucleic acid fragments of the instant invention may be used to isolate cDNAs and genes encoding homologous proteins from the same or other plant species. Isolation of homologous genes using sequence-dependent protocols is well known in the art. Examples of sequence-dependent protocols include, but are not limited to, methods of nucleic acid hybridization, and methods of DNA and RNA amplification as exemplified by various uses of nucleic acid amplification technologies (e.g., polymerase chain reaction, ligase chain reaction).

For example, genes encoding other Myb-related transcription factors, either as cDNAs or genomic DNAs, could be isolated directly by using all or a portion of the instant nucleic acid fragments as DNA hybridization probes to screen libraries from any desired plant employing methodology well known to those skilled in the art. Specific oligonucleotide probes based upon the instant nucleic acid sequences can be designed and synthesized by methods known in the art (Maniatis). Moreover, the entire sequences can be used directly to synthesize DNA probes by methods known to the skilled artisan such as random primer DNA labeling, nick translation, or end-labeling techniques, or RNA probes using available *in vitro* transcription systems. In addition, specific primers can be designed and used to amplify a part or all of the instant sequences. The resulting amplification products can be labeled directly during amplification reactions or labeled after amplification reactions, and used as probes to isolate full length cDNA or genomic fragments under conditions of appropriate stringency.

In addition, two short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols to amplify longer nucleic acid fragments encoding homologous genes from DNA or RNA. The polymerase chain reaction may also be performed on a library of cloned nucleic acid fragments wherein the sequence of one primer is derived from the instant nucleic acid fragments, and the sequence of the other primer takes advantage of the presence of the polyadenylic acid tracts to the 3' end of the mRNA precursor encoding plant genes. Alternatively, the second primer sequence may be based upon sequences derived from the cloning vector. For example, the skilled artisan can follow the RACE protocol (Frohman et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:8998-9002) to generate cDNAs by using PCR to amplify copies of the region between a single point in the transcript and the 3' or 5' end. Primers oriented in the 3' and 5' directions can be designed from the instant sequences. Using commercially available 3' RACE or 5' RACE systems (BRL), specific 3' or 5' cDNA fragments can be isolated (Ohara et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:5673-5677; Loh et al. (1989) *Science* 243:217-220). Products generated by the 3' and 5' RACE procedures can be combined to generate full-length cDNAs (Frohman

and Martin (1989) *Techniques* 1:165). Consequently, a polynucleotide comprising a nucleotide sequence of at least one of 60 (preferably one of at least 40, most preferably one of at least 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61 and the complement of such nucleotide sequences may be used in such methods to obtain a nucleic acid fragment encoding a substantial portion of an amino acid sequence of a polypeptide (such as a Myb-related transcription factor. The present invention relates to a method of obtaining a nucleic acid fragment encoding a substantial portion of a polypeptide of a gene (such as Myb-related transcription factor) preferably a substantial portion of a plant polypeptide of a gene, comprising the steps of: synthesizing an oligonucleotide primer comprising a nucleotide sequence of at least one of 60 (preferably at least one of 40, most preferably at least one of 30) contiguous nucleotides derived from a nucleotide sequence selected from the group consisting of SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, 45, 47, 49, 51, 53, 55, 57, 59, and 61 and the complement of such nucleotide sequences; and amplifying a nucleic acid fragment (preferably a cDNA inserted in a cloning vector) using the oligonucleotide primer. The amplified nucleic acid fragment preferably will encode a portion of a polypeptide.

Availability of the instant nucleotide and deduced amino acid sequences facilitates immunological screening of cDNA expression libraries. Synthetic peptides representing portions of the instant amino acid sequences may be synthesized. These peptides can be used to immunize animals to produce polyclonal or monoclonal antibodies with specificity for peptides or proteins comprising the amino acid sequences. These antibodies can be then be used to screen cDNA expression libraries to isolate full-length cDNA clones of interest (Lerner (1984) *Adv. Immunol.* 36:1-34; Maniatis).

The nucleic acid fragments of the instant invention may be used to create transgenic plants in which the disclosed polypeptides are present at higher or lower levels than normal or in cell types or developmental stages in which they are not normally found. This would have the effect of altering the level of expression of Myb-regulated genes in those cells, and consequently the phenotype affected by those Myb-regulated genes.

Overexpression of the proteins of the instant invention may be accomplished by first constructing a chimeric gene in which the coding region is operably linked to a promoter capable of directing expression of a gene in the desired tissues at the desired stage of development. For reasons of convenience, the chimeric gene may comprise promoter sequences and translation leader sequences derived from the same genes. 3' Non-coding sequences encoding transcription termination signals may also be provided. The instant chimeric gene may also comprise one or more introns in order to facilitate gene expression.

Plasmid vectors comprising the instant chimeric gene can then be constructed. The choice of plasmid vector is dependent upon the method that will be used to transform host plants. The skilled artisan is well aware of the genetic elements that must be present on the plasmid vector in order to successfully transform, select and propagate host cells containing the chimeric gene. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al. (1985) *EMBO J.* 4:2411-2418; De Almeida et al. (1989) *Mol. Gen. Genetics* 218:78-86), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

For some applications it may be useful to direct the instant polypeptides to different cellular compartments, or to facilitate its secretion from the cell. It is thus envisioned that the chimeric gene described above may be further supplemented by altering the coding sequence to encode the instant polypeptides with appropriate intracellular targeting sequences such as transit sequences (Keegstra (1989) *Cell* 56:247-253), signal sequences or sequences encoding endoplasmic reticulum localization (Chrispeels (1991) *Ann. Rev. Plant Phys. Plant Mol. Biol.* 42:21-53), or nuclear localization signals (Raikhel (1992) *Plant Phys.* 100:1627-1632) added and/or with targeting sequences that are already present removed. While the references cited give examples of each of these, the list is not exhaustive and more targeting signals of utility may be discovered in the future.

It may also be desirable to reduce or eliminate expression of genes encoding the instant polypeptides in plants for some applications. In order to accomplish this, a chimeric gene designed for co-suppression of the instant polypeptide can be constructed by linking a gene or gene fragment encoding that polypeptide to plant promoter sequences. Alternatively, a chimeric gene designed to express antisense RNA for all or part of the instant nucleic acid fragment can be constructed by linking the gene or gene fragment in reverse orientation to plant promoter sequences. Either the co-suppression or antisense chimeric genes could be introduced into plants via transformation wherein expression of the corresponding endogenous genes are reduced or eliminated.

Molecular genetic solutions to the generation of plants with altered gene expression have a decided advantage over more traditional plant breeding approaches. Changes in plant phenotypes can be produced by specifically inhibiting expression of one or more genes by antisense inhibition or cosuppression (U.S. Patent Nos. 5,190,931, 5,107,065 and 5,283,323). An antisense or cosuppression construct would act as a dominant negative regulator of gene activity. While conventional mutations can yield negative regulation of gene activity these effects are most likely recessive. The dominant negative regulation available with a transgenic approach may be advantageous from a breeding perspective. In

addition, the ability to restrict the expression of specific phenotype to the reproductive tissues of the plant by the use of tissue specific promoters may confer agronomic advantages relative to conventional mutations which may have an effect in all tissues in which a mutant gene is ordinarily expressed.

5 The person skilled in the art will know that special considerations are associated with the use of antisense or cosuppression technologies in order to reduce expression of particular genes. For example, the proper level of expression of sense or antisense genes may require the use of different chimeric genes utilizing different regulatory elements known to the skilled artisan. Once transgenic plants are obtained by one of the methods described above,
10 it will be necessary to screen individual transgenics for those that most effectively display the desired phenotype. Accordingly, the skilled artisan will develop methods for screening large numbers of transformants. The nature of these screens will generally be chosen on practical grounds, and is not an inherent part of the invention. For example, one can screen by looking for changes in gene expression by using antibodies specific for the protein
15 encoded by the gene being suppressed, or one could establish assays that specifically measure enzyme activity. A preferred method will be one which allows large numbers of samples to be processed rapidly, since it will be expected that a large number of transformants will be negative for the desired phenotype.

The instant polypeptides (or portions thereof) may be produced in heterologous host
20 cells, particularly in the cells of microbial hosts, and can be used to prepare antibodies to the these proteins by methods well known to those skilled in the art. The antibodies are useful for detecting the polypeptides of the instant invention *in situ* in cells or *in vitro* in cell extracts. Preferred heterologous host cells for production of the instant polypeptides are microbial hosts. Microbial expression systems and expression vectors containing regulatory
25 sequences that direct high level expression of foreign proteins are well known to those skilled in the art. Any of these could be used to construct a chimeric gene for production of the instant polypeptides. This chimeric gene could then be introduced into appropriate microorganisms via transformation to provide high level expression of the encoded Myb-related transcription factors. An example of a vector for high level expression of the
30 instant polypeptides in a bacterial host is provided (Example 6).

All or a substantial portion of the nucleic acid fragments of the instant invention may also be used as probes for genetically and physically mapping the genes that they are a part of, and as markers for traits linked to those genes. Such information may be useful in plant breeding in order to develop lines with desired phenotypes. For example, the instant nucleic
35 acid fragments may be used as restriction fragment length polymorphism (RFLP) markers. Southern blots (Maniatis) of restriction-digested plant genomic DNA may be probed with the nucleic acid fragments of the instant invention. The resulting banding patterns may then be subjected to genetic analyses using computer programs such as MapMaker (Lander et al.

(1987) *Genomics* 1:174-181) in order to construct a genetic map. In addition, the nucleic acid fragments of the instant invention may be used to probe Southern blots containing restriction endonuclease-treated genomic DNAs of a set of individuals representing parent and progeny of a defined genetic cross. Segregation of the DNA polymorphisms is noted and used to calculate the position of the instant nucleic acid sequence in the genetic map previously obtained using this population (Botstein et al. (1980) *Am. J. Hum. Genet.* 32:314-331).

The production and use of plant gene-derived probes for use in genetic mapping is described in Bernatzky and Tanksley (1986) *Plant Mol. Biol. Reporter* 4:37-41. Numerous publications describe genetic mapping of specific cDNA clones using the methodology outlined above or variations thereof. For example, F2 intercross populations, backcross populations, randomly mated populations, near isogenic lines, and other sets of individuals may be used for mapping. Such methodologies are well known to those skilled in the art.

Nucleic acid probes derived from the instant nucleic acid sequences may also be used for physical mapping (i.e., placement of sequences on physical maps; see Hoheisel et al. In: *Nonmammalian Genomic Analysis: A Practical Guide*, Academic press 1996, pp. 319-346, and references cited therein).

In another embodiment, nucleic acid probes derived from the instant nucleic acid sequences may be used in direct fluorescence *in situ* hybridization (FISH) mapping (Trask (1991) *Trends Genet.* 7:149-154). Although current methods of FISH mapping favor use of large clones (several to several hundred KB; see Laan et al. (1995) *Genome Res.* 5:13-20), improvements in sensitivity may allow performance of FISH mapping using shorter probes.

A variety of nucleic acid amplification-based methods of genetic and physical mapping may be carried out using the instant nucleic acid sequences. Examples include allele-specific amplification (Kazazian (1989) *J. Lab. Clin. Med.* 11:95-96), polymorphism of PCR-amplified fragments (CAPS; Sheffield et al. (1993) *Genomics* 16:325-332), allele-specific ligation (Landegren et al. (1988) *Science* 241:1077-1080), nucleotide extension reactions (Sokolov (1990) *Nucleic Acid Res.* 18:3671), Radiation Hybrid Mapping (Walter et al. (1997) *Nat. Genet.* 7:22-28) and Happy Mapping (Dear and Cook (1989) *Nucleic Acid Res.* 17:6795-6807). For these methods, the sequence of a nucleic acid fragment is used to design and produce primer pairs for use in the amplification reaction or in primer extension reactions. The design of such primers is well known to those skilled in the art. In methods employing PCR-based genetic mapping, it may be necessary to identify DNA sequence differences between the parents of the mapping cross in the region corresponding to the instant nucleic acid sequence. This, however, is generally not necessary for mapping methods.

Loss of function mutant phenotypes may be identified for the instant cDNA clones either by targeted gene disruption protocols or by identifying specific mutants for these

genes contained in a maize population carrying mutations in all possible genes (Ballinger and Benzer (1989) *Proc. Natl. Acad. Sci USA* 86:9402-9406; Koes et al. (1995) *Proc. Natl. Acad. Sci USA* 92:8149-8153; Bensen et al. (1995) *Plant Cell* 7:75-84). The latter approach may be accomplished in two ways. First, short segments of the instant nucleic acid fragments may be used in polymerase chain reaction protocols in conjunction with a mutation tag sequence primer on DNAs prepared from a population of plants in which Mutator transposons or some other mutation-causing DNA element has been introduced (see Bensen, *supra*). The amplification of a specific DNA fragment with these primers indicates the insertion of the mutation tag element in or near the plant gene encoding the instant polypeptide. Alternatively, the instant nucleic acid fragment may be used as a hybridization probe against PCR amplification products generated from the mutation population using the mutation tag sequence primer in conjunction with an arbitrary genomic site primer, such as that for a restriction enzyme site-anchored synthetic adaptor. With either method, a plant containing a mutation in the endogenous gene encoding the instant polypeptide can be identified and obtained. This mutant plant can then be used to determine or confirm the natural function of the instant polypeptides disclosed herein.

EXAMPLES

The present invention is further defined in the following Examples, in which all parts and percentages are by weight and degrees are Celsius, unless otherwise stated. It should be understood that these Examples, while indicating preferred embodiments of the invention, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can ascertain the essential characteristics of this invention, and without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

EXAMPLE 1

Composition of cDNA Libraries: Isolation and Sequencing of cDNA Clones

cDNA libraries representing mRNAs from various corn, rice, soybean and wheat tissues were prepared. The characteristics of the libraries are described below.

TABLE 2
cDNA Libraries from Corn, Rice, Soybean and Wheat

Library	Tissue	Clone
chpc8	Corn (<i>Zea mays</i> L.) (MBS847) 8 Day Old Shoot Treated 8 Hours With PDO Herbicide MK593*	chpc8.pk0002.d9
csi1n	Corn (<i>Zea mays</i> L.) Silk**	csi1n.pk0008.b5 csi1n.pk0028.h7
cta1n	Corn (<i>Zea mays</i> L.) Tassel**	cta1n.pk0079.e9
p0008	Corn (<i>Zea mays</i> L.) 3 Week Old Leaf	p0008.cb31d06r
p0026	Corn (<i>Zea mays</i> L.) Regenerating Callus (Hi-II 223a and 1129e), 5 Days After Auxin Removal	p0026.ccrbd36rb
rds3c	Rice (<i>Oryza sativa</i>) Developing Seed From Top of the Plant	rds3c.pk002.c6
rl0n	Rice (<i>Oryza sativa</i>) 15 Day Old Leaf**	rl0n.pk082.c13
rlr24	Resistant Rice (<i>Oryza sativa</i>) Leaf 15 Days After Germination, 24 Hours After Infection of Strain <i>Magnaporthe grisea</i> 4360-R-62 (AVR2-YAMO)	rlr24.pk0090.f5
rlr48	Resistant Rice (<i>Oryza sativa</i>) Leaf 15 Days After Germination, 48 Hours After Infection of Strain <i>Magnaporthe grisea</i> 4360-R-62 (AVR2-YAMO)	rlr48.pk0012.c11
rr1	Rice (<i>Oryza sativa</i>) Root of Two Week Old Developing Seedling	rr1.pk0027.g9 rr1.pk0037.g7 rr1.pk077.n9 rr1.pk088.p6
se6	Soybean (<i>Glycine max</i> L) Embryo, 26 Days After Flowering	se6.pk0048.a12
ses9c	Soybean (<i>Glycine max</i> L) Embryogenic Suspension	ses9c.pk002.o16
sfl1	Soybean (<i>Glycine max</i> L) Immature Flower	sfl1.pk0003.a3 sfl1.pk0032.g4 sfl1.pk0086.a9 sfl1.pk0091.a2 sfl1.pk0105.e6 sfl1.pk125.p19
sgs4c	Soybean (<i>Glycine max</i> L) Seed 2 Days After Germination	sgs4c.pk004.j24
sl2	Soybean (<i>Glycine max</i> L) Two-Week-Old Developing Seedling Treated With 2.5 ppm chlorimuron	sl2.pk127.e14
src3c	Soybean (<i>Glycine max</i> L) 8 Day Old Root Infected With Cyst Nematode	src3c.pk010.i22
srr3c	Soybean (<i>Glycine max</i> L) 8 Day Old Root	srr3c.pk002.k6
wdk3c	Wheat (<i>Triticum aestivum</i> L) Developing Kernel, 14 Days After Anthesis	wdk3c.pk006.n12
wlm1	Wheat (<i>Triticum aestivum</i> L) Seedling 1 Hour After Inoculation With <i>Erysiphe graminis</i> f. <i>sp tritici</i>	wlm1.pk0027.a5
wr1	Wheat (<i>Triticum aestivum</i> L) Root From 7 Day Old Seedling Light Grown	wr1.pk0139.g11

*Application of 2-[(2,4-dihydro-2,6,9-trimethyl[1]benzothioapyrano[4,3-c]pyrazol-8-

yl)carbonyl]-1,3-cyclohexanedione *S,S*-dioxide; synthesis and methods of using this compound are described in WO 97/19087, incorporated herein by reference.

**These libraries were normalized essentially as described in U.S. Patent No. 5,482,845, incorporated herein by reference.

cDNA libraries may be prepared by any one of many methods available. For example, the cDNAs may be introduced into plasmid vectors by first preparing the cDNA libraries in Uni-ZAP™ XR vectors according to the manufacturer's protocol (Stratagene Cloning Systems, La Jolla, CA). The Uni-ZAP™ XR libraries are converted into plasmid libraries according to the protocol provided by Stratagene. Upon conversion, cDNA inserts will be contained in the plasmid vector pBluescript. In addition, the cDNAs may be introduced directly into precut Bluescript II SK(+) vectors (Stratagene) using T4 DNA ligase (New England Biolabs), followed by transfection into DH10B cells according to the manufacturer's protocol (GIBCO BRL Products). Once the cDNA inserts are in plasmid vectors, plasmid DNAs are prepared from randomly picked bacterial colonies containing recombinant pBluescript plasmids, or the insert cDNA sequences are amplified via polymerase chain reaction using primers specific for vector sequences flanking the inserted cDNA sequences. Amplified insert DNAs or plasmid DNAs are sequenced in dye-primer sequencing reactions to generate partial cDNA sequences (expressed sequence tags or "ESTs"; see Adams et al., (1991) *Science* 252:1651-1656). The resulting ESTs are analyzed using a Perkin Elmer Model 377 fluorescent sequencer.

Determination of complete nucleotide sequence of cDNA inserts may be accomplished by a number of methods well-known to those skilled in the art (Maniatis). For example, this may be accomplished stepwise, wherein oligonucleotides near the 5' or 3' end of the sequence may be synthesized which can then serve as primers for sequencing reactions that will extend the known sequence. Another set of oligonucleotides near the 5' or 3' end of the new sequence in the next round prime another set of sequencing reactions to obtain more sequence information. These steps are repeated until the complete nucleotide sequence is determined.

EXAMPLE 2

Identification of cDNA Clones

cDNA clones encoding Myb-related transcription factors were identified by conducting BLAST (Basic Local Alignment Search Tool; Altschul et al. (1993) *J. Mol. Biol.* 215:403-410; see also www.ncbi.nlm.nih.gov/BLAST/) searches for similarity to sequences contained in the BLAST "nr" database (comprising all non-redundant GenBank CDS translations, sequences derived from the 3-dimensional structure Brookhaven Protein Data Bank, the last major release of the SWISS-PROT protein sequence database, EMBL, and DDBJ databases). The cDNA sequences obtained in Example 1 were analyzed for similarity to all publicly available DNA sequences contained in the "nr" database using the BLASTN

algorithm provided by the National Center for Biotechnology Information (NCBI). The DNA sequences were translated in all reading frames and compared for similarity to all publicly available protein sequences contained in the "nr" database using the BLASTX algorithm (Gish and States (1993) *Nat. Genet.* 3:266-272) provided by the NCBI. For convenience, the P-value (probability) of observing a match of a cDNA sequence to a sequence contained in the searched databases merely by chance as calculated by BLAST are reported herein as "pLog" values, which represent the negative of the logarithm of the reported P-value. Accordingly, the greater the pLog value, the greater the likelihood that the cDNA sequence and the BLAST "hit" represent homologous proteins.

EXAMPLE 3

Characterization of cDNA Clones Encoding Myb-related Transcription Factors

The BLASTX search using the EST sequence from clone cta1n.pk0079.e9 revealed similarity of the protein encoded by the cDNA to a Myb-related transcription factor from *Craterostigma plantagineum* (NCBI Identifier No. gi 1002800). The BLASTX search using the EST sequence from clone rr1.pk077.n9 revealed similarity of the protein encoded by the cDNA to a Myb-related transcription factor from *Arabidopsis thaliana* (NCBI Identifier No. gi 1732513). The BLASTX search using the EST sequence from clone sfl1.pk0032.g4 revealed similarity of the protein encoded by the cDNA to a Myb-related transcription factor from *Pisum sativum* (NCBI Identifier No. gi 1841475). The BLASTX search using the EST sequence from clone wr1.pk0139.g11 revealed similarity of the protein encoded by the cDNA to a Myb-related transcription factor from *Arabidopsis thaliana* (NCBI Identifier No. gi 2832500). The BLAST results for each of these ESTs are shown in Table 3:

TABLE 3

BLAST Results for Clones Encoding Polypeptides Homologous to Plant Myb-related Transcription Factors

Clone	BLAST pLog Score
cta1n.pk0079.e9	39.00
rr1.pk077.n9	27.70
sfl1.pk0032.g4	38.50
wr1.pk0139.g11	16.00

The sequence of a portion of the cDNA insert from clone cta1n.pk0079.e9 is shown in SEQ ID NO:7; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:8. The sequence of a portion of the cDNA insert from clone rr1.pk077.n9 is shown in SEQ ID NO:11; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:12. The sequence of a portion of the cDNA insert from clone sfl1.pk0032.g4 is shown in SEQ ID NO:27; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:28. The

sequence of a portion of the cDNA insert from clone wr1.pk0139.g11 is shown in SEQ ID NO:55; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:56. BLAST scores and probabilities indicate that the instant nucleic acid fragments encode portions of a Myb-related transcription factor protein.

The BLASTX search using the EST sequences from clones rl0n.pk082.c13 and sgs4c.pk004.j24 revealed similarity of the proteins encoded by the cDNAs to a Myb-related transcription factor protein from *Pisum sativum* (NCBI Identifier No. gi 82307). The BLAST results for each of these ESTs are shown in Table 4:

TABLE 4

BLAST Results for Clones Encoding Polypeptides Homologous to Plant Myb-related Transcription Factors

Clone	BLAST pLog Score
rl0n.pk082.c13	62.50
sgs4c.pk004.j24	47.50

The sequence of a portion of the cDNA insert from clone rl0n.pk082.c13 is shown in SEQ ID NO:23; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:24. The sequence of a portion of the cDNA insert from clone sgs4c.pk004.j24 is shown in SEQ ID NO:51; the deduced amino acid sequence of this cDNA is shown in SEQ ID NO:52. BLAST scores and probabilities indicate that the instant nucleic acid fragments encode portions of a Myb-related transcription factor protein.

The BLASTX search using the sequences from clones listed in Table 5 revealed similarity of the polypeptides encoded by the cDNAs to Myb-related transcription factors from different plant species including *Pisum sativum* (NCBI General Identification No. 1841475), *Arabidopsis thaliana* (NCBI General Identification Nos. 3941480 and 3941528), *Craterostigma plantagineum* (NCBI General Identification Nos. 1002796, 1002798, and 1002800), and *Antirrhinum majus* (NCBI General Identification No. 82307). Shown in Table 5 are the BLAST results for individual ESTs ("EST"), the sequences of the entire cDNA inserts comprising the indicated cDNA clones ("FIS"), contigs assembled from two or more ESTs ("Contig"), contigs assembled from an FIS and one or more ESTs ("Contig*"), or sequences encoding the entire protein derived from an FIS, a contig, or an FIS and PCR ("CGS"):

TABLE 5
BLAST Results for Sequences Encoding Polypeptides Homologous
to Myb-related Transcription Factors

Clone	Status	NCBI General Identification No.	pLog Score
Contig of: csi1n.pk0008.b5 csi1n.pk0028.h7	Contig	1841475	54.40
Contig of: p0008.cb3ld06r p0026.ccrbd36rb	Contig	3941480	70.10
chpc8.pk0002.d9	EST	1002800	44.70
cta1n.pk0079.e9	CGS	1002800	68.52
Contig of: rr1.pk0027.g9 rr1.pk077.n9	CGS	3941480	50.00
rr1.pk088.p6	EST	3941480	22.52
rr1.pk0037.g7	EST	1002798	79.70
rds3c.pk002.c6	EST	1002798	62.00
Contig of: rlr24.pk0090.f5 rlr48.pk0012.c11	Contig	1002800	55.04
rl0n.pk082.c13	CGS	82307	84.22
sfl1.pk0032.g4	CGS	1841475	96.52
sfl1.pk0086.a9	CGS	1841475	92.22
sfl1.pk0091.a2	CGS	1841475	75.10
sfl1.pk0105.e6	CGS	1841475	96.40
sfl1.pk125.p19	FIS	1841475	90.52
se6.pk0048.a12	FIS	1002798	66.70
sfl1.pk0003.a3	CGS	1002796	59.70
srr3c.pk002.k6	CGS	1002798	77.40
ses9c.pk002.o16	CGS	1002798	73.70
sl2.pk127.e14	CGS	1002800	60.70
src3c.pk010.i22	CGS	1002800	57.05
sgs4c.pk004.j24	CGS	82307	90.52
wr1.pk0139.g11	FIS	3941480	48.52
wdk3c.pk006.n12	EST	1002796	43.70
wlm1.pk0027.a5	EST	3941528	73.70

5 Figure 1 presents an alignment of the amino acid sequences set forth in SEQ ID
NOs:10, 14, 26, 30, 32, 34, 36, 42, 44, 46, 48, 50, and 54 and the *Pisum sativum* sequence

(NCBI General Identification No. 1841475; SEQ ID NO:63). The data in Table 6 represents a calculation of the percent identity of the amino acid sequences set forth in SEQ ID NOS:10, 14, 26, 30, 32, 34, 36, 42, 44, 46, 48, 50, and 54 and the *Pisum sativum* sequence (NCBI General Identification No. 1841475; SEQ ID NO:63).

TABLE 6

Percent Identity of Amino Acid Sequences Deduced From the Nucleotide Sequences of cDNA Clones Encoding Polypeptides Homologous to Myb-related Transcription Factors

SEQ ID NO.	Percent Identity to NCBI General Identification No. 1841475
10	43.8
14	33.2
26	30.9
30	75.6
32	74.1
34	64.1
36	77.7
42	43.8
44	48.5
46	46.5
48	45.6
50	46.1
54	31.8

- 10 Sequence alignments and percent identity calculations were performed using the Megalign program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Multiple alignment of the sequences was performed using the Clustal method of alignment (Higgins and Sharp (1989) *CABIOS* 5:151-153) with the default parameters (GAP PENALTY=10, GAP LENGTH PENALTY=10). Default parameters for
- 15 pairwise alignments using the Clustal method were KTUPLE 1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. Sequence alignments and BLAST scores and probabilities indicate that the nucleic acid fragments comprising the instant cDNA clones encode all or a substantial portion of a Myb-related transcription factor. These sequences represent the first soybean and wheat sequences encoding Myb-related transcription factors.
- 20 Nucleic acid fragments encoding Myb-related transcription factors have previously been isolated from rice and corn (Marocco et al. (1989) *Mol Gen Genet* 216:183-187; Pandolfi et al. (1997) *Plant Physiol* 114:747).

EXAMPLE 4

Expression of Chimeric Genes in Monocot Cells

A chimeric gene comprising a cDNA encoding the instant polypeptide in sense orientation with respect to the maize 27 kD zein promoter that is located 5' to the cDNA fragment, and the 10 kD zein 3' end that is located 3' to the cDNA fragment, can be constructed. The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites (NcoI or SmaI) can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the digested vector pML103 as described below.

Amplification is then performed in a standard PCR. The amplified DNA is then digested with restriction enzymes NcoI and SmaI and fractionated on an agarose gel. The appropriate band can be isolated from the gel and combined with a 4.9 kb NcoI-SmaI fragment of the plasmid pML103. Plasmid pML103 has been deposited under the terms of the Budapest Treaty at ATCC (American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209), and bears accession number ATCC 97366. The DNA segment from pML103 contains a 1.05 kb Sall-NcoI promoter fragment of the maize 27 kD zein gene and a 0.96 kb SmaI-SalI fragment from the 3' end of the maize 10 kD zein gene in the vector pGem9Zf(+) (Promega). Vector and insert DNA can be ligated at 15°C overnight, essentially as described (Maniatis). The ligated DNA may then be used to transform *E. coli* XL1-Blue (Epicurian Coli XL-1 Blue™; Stratagene). Bacterial transformants can be screened by restriction enzyme digestion of plasmid DNA and limited nucleotide sequence analysis using the dideoxy chain termination method (Sequenase™ DNA Sequencing Kit; U.S. Biochemical). The resulting plasmid construct would comprise a chimeric gene encoding, in the 5' to 3' direction, the maize 27 kD zein promoter, a cDNA fragment encoding the instant polypeptide, and the 10 kD zein 3' region.

The chimeric gene described above can then be introduced into corn cells by the following procedure. Immature corn embryos can be dissected from developing caryopses derived from crosses of the inbred corn lines H99 and LH132. The embryos are isolated 10 to 11 days after pollination when they are 1.0 to 1.5 mm long. The embryos are then placed with the axis-side facing down and in contact with agarose-solidified N6 medium (Chu et al. (1975) *Sci. Sin. Peking* 18:659-668). The embryos are kept in the dark at 27°C. Friable embryogenic callus consisting of undifferentiated masses of cells with somatic proembryoids and embryoids borne on suspensor structures proliferates from the scutellum of these immature embryos. The embryogenic callus isolated from the primary explant can be cultured on N6 medium and sub-cultured on this medium every 2 to 3 weeks.

The plasmid, p35S/Ac (obtained from Dr. Peter Eckes, Hoechst Ag, Frankfurt, Germany) may be used in transformation experiments in order to provide for a selectable marker. This plasmid contains the *Pat* gene (see European Patent Publication 0 242 236)

which encodes phosphinothricin acetyl transferase (PAT). The enzyme PAT confers resistance to herbicidal glutamine synthetase inhibitors such as phosphinothricin. The *pat* gene in p35S/Ac is under the control of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*.

The particle bombardment method (Klein et al. (1987) *Nature* 327:70-73) may be used to transfer genes to the callus culture cells. According to this method, gold particles (1 μ m in diameter) are coated with DNA using the following technique. Ten μ g of plasmid DNAs are added to 50 μ L of a suspension of gold particles (60 mg per mL). Calcium chloride (50 μ L of a 2.5 M solution) and spermidine free base (20 μ L of a 1.0 M solution) are added to the particles. The suspension is vortexed during the addition of these solutions. After 10 minutes, the tubes are briefly centrifuged (5 sec at 15,000 rpm) and the supernatant removed. The particles are resuspended in 200 μ L of absolute ethanol, centrifuged again and the supernatant removed. The ethanol rinse is performed again and the particles resuspended in a final volume of 30 μ L of ethanol. An aliquot (5 μ L) of the DNA-coated gold particles can be placed in the center of a Kapton™ flying disc (Bio-Rad Labs). The particles are then accelerated into the corn tissue with a Biolistic™ PDS-1000/He (Bio-Rad Instruments, Hercules CA), using a helium pressure of 1000 psi, a gap distance of 0.5 cm and a flying distance of 1.0 cm.

For bombardment, the embryogenic tissue is placed on filter paper over agarose-solidified N6 medium. The tissue is arranged as a thin lawn and covered a circular area of about 5 cm in diameter. The petri dish containing the tissue can be placed in the chamber of the PDS-1000/He approximately 8 cm from the stopping screen. The air in the chamber is then evacuated to a vacuum of 28 inches of Hg. The macrocarrier is accelerated with a helium shock wave using a rupture membrane that bursts when the He pressure in the shock tube reaches 1000 psi.

Seven days after bombardment the tissue can be transferred to N6 medium that contains glufosinate (2 mg per liter) and lacks casein or proline. The tissue continues to grow slowly on this medium. After an additional 2 weeks the tissue can be transferred to fresh N6 medium containing glufosinate. After 6 weeks, areas of about 1 cm in diameter of actively growing callus can be identified on some of the plates containing the glufosinate-supplemented medium. These calli may continue to grow when sub-cultured on the selective medium.

Plants can be regenerated from the transgenic callus by first transferring clusters of tissue to N6 medium supplemented with 0.2 mg per liter of 2,4-D. After two weeks the tissue can be transferred to regeneration medium (Fromm et al. (1990) *Bio/Technology* 8:833-839).

EXAMPLE 5

Expression of Chimeric Genes in Dicot Cells

A seed-specific expression cassette composed of the promoter and transcription terminator from the gene encoding the β subunit of the seed storage protein phaseolin from the bean *Phaseolus vulgaris* (Doyle et al. (1986) *J. Biol. Chem.* 261:9228-9238) can be used for expression of the instant polypeptides in transformed soybean. The phaseolin cassette includes about 500 nucleotides upstream (5') from the translation initiation codon and about 1650 nucleotides downstream (3') from the translation stop codon of phaseolin. Between the 5' and 3' regions are the unique restriction endonuclease sites Nco I (which includes the ATG translation initiation codon), Sma I, Kpn I and Xba I. The entire cassette is flanked by Hind III sites.

The cDNA fragment of this gene may be generated by polymerase chain reaction (PCR) of the cDNA clone using appropriate oligonucleotide primers. Cloning sites can be incorporated into the oligonucleotides to provide proper orientation of the DNA fragment when inserted into the expression vector. Amplification is then performed as described above, and the isolated fragment is inserted into a pUC18 vector carrying the seed expression cassette.

Soybean embryos may then be transformed with the expression vector comprising sequences encoding the instant polypeptides. To induce somatic embryos, cotyledons, 3-5 mm in length dissected from surface sterilized, immature seeds of the soybean cultivar A2872, can be cultured in the light or dark at 26°C on an appropriate agar medium for 6-10 weeks. Somatic embryos which produce secondary embryos are then excised and placed into a suitable liquid medium. After repeated selection for clusters of somatic embryos which multiplied as early, globular staged embryos, the suspensions are maintained as described below.

Soybean embryogenic suspension cultures can be maintained in 35 mL liquid media on a rotary shaker, 150 rpm, at 26°C with florescent lights on a 16:8 hour day/night schedule. Cultures are subcultured every two weeks by inoculating approximately 35 mg of tissue into 35 mL of liquid medium.

Soybean embryogenic suspension cultures may then be transformed by the method of particle gun bombardment (Klein et al. (1987) *Nature* (London) 327:70-73, U.S. Patent No. 4,945,050). A DuPont Biolistic™ PDS1000/HE instrument (helium retrofit) can be used for these transformations.

A selectable marker gene which can be used to facilitate soybean transformation is a chimeric gene composed of the 35S promoter from Cauliflower Mosaic Virus (Odell et al. (1985) *Nature* 313:810-812), the hygromycin phosphotransferase gene from plasmid pJR225 (from *E. coli*; Gritz et al. (1983) *Gene* 25:179-188) and the 3' region of the nopaline synthase gene from the T-DNA of the Ti plasmid of *Agrobacterium tumefaciens*. The seed expression

cassette comprising the phaseolin 5' region, the fragment encoding the instant polypeptide and the phaseolin 3' region can be isolated as a restriction fragment. This fragment can then be inserted into a unique restriction site of the vector carrying the marker gene.

To 50 μ L of a 60 mg/mL 1 μ m gold particle suspension is added (in order): 5 μ L DNA (1 μ g/ μ L), 20 μ L spermidine (0.1 M), and 50 μ L CaCl_2 (2.5 M). The particle preparation is then agitated for three minutes, spun in a microfuge for 10 seconds and the supernatant removed. The DNA-coated particles are then washed once in 400 μ L 70% ethanol and resuspended in 40 μ L of anhydrous ethanol. The DNA/particle suspension can be sonicated three times for one second each. Five μ L of the DNA-coated gold particles are then loaded on each macro carrier disk.

Approximately 300-400 mg of a two-week-old suspension culture is placed in an empty 60x15 mm petri dish and the residual liquid removed from the tissue with a pipette. For each transformation experiment, approximately 5-10 plates of tissue are normally bombarded. Membrane rupture pressure is set at 1100 psi and the chamber is evacuated to a vacuum of 28 inches mercury. The tissue is placed approximately 3.5 inches away from the retaining screen and bombarded three times. Following bombardment, the tissue can be divided in half and placed back into liquid and cultured as described above.

Five to seven days post bombardment, the liquid media may be exchanged with fresh media, and eleven to twelve days post bombardment with fresh media containing 50 mg/mL hygromycin. This selective media can be refreshed weekly. Seven to eight weeks post bombardment, green, transformed tissue may be observed growing from untransformed, necrotic embryogenic clusters. Isolated green tissue is removed and inoculated into individual flasks to generate new, clonally propagated, transformed embryogenic suspension cultures. Each new line may be treated as an independent transformation event. These suspensions can then be subcultured and maintained as clusters of immature embryos or regenerated into whole plants by maturation and germination of individual somatic embryos.

EXAMPLE 6

Expression of Chimeric Genes in Microbial Cells

The cDNAs encoding the instant polypeptides can be inserted into the T7 *E. coli* expression vector pBT430. This vector is a derivative of pET-3a (Rosenberg et al. (1987) *Gene* 56:125-135) which employs the bacteriophage T7 RNA polymerase/T7 promoter system. Plasmid pBT430 was constructed by first destroying the EcoR I and Hind III sites in pET-3a at their original positions. An oligonucleotide adaptor containing EcoR I and Hind III sites was inserted at the BamH I site of pET-3a. This created pET-3aM with additional unique cloning sites for insertion of genes into the expression vector. Then, the Nde I site at the position of translation initiation was converted to an Nco I site using oligonucleotide-directed mutagenesis. The DNA sequence of pET-3aM in this region, 5'-CATATGG, was converted to 5'-CCCATGG in pBT430.

Plasmid DNA containing a cDNA may be appropriately digested to release a nucleic acid fragment encoding the protein. This fragment may then be purified on a 1% NuSieve GTG™ low melting agarose gel (FMC). Buffer and agarose contain 10 µg/ml ethidium bromide for visualization of the DNA fragment. The fragment can then be purified from the agarose gel by digestion with GELase™ (Epicentre Technologies) according to the manufacturer's instructions, ethanol precipitated, dried and resuspended in 20 µL of water. Appropriate oligonucleotide adapters may be ligated to the fragment using T4 DNA ligase (New England Biolabs, Beverly, MA). The fragment containing the ligated adapters can be purified from the excess adapters using low melting agarose as described above. The vector pBT430 is digested, dephosphorylated with alkaline phosphatase (NEB) and deproteinized with phenol/chloroform as described above. The prepared vector pBT430 and fragment can then be ligated at 16°C for 15 hours followed by transformation into DH5 electrocompetent cells (GIBCO BRL). Transformants can be selected on agar plates containing LB media and 100 µg/mL ampicillin. Transformants containing the gene encoding the instant polypeptide are then screened for the correct orientation with respect to the T7 promoter by restriction enzyme analysis.

For high level expression, a plasmid clone with the cDNA insert in the correct orientation relative to the T7 promoter can be transformed into *E. coli* strain BL21(DE3) (Studier et al. (1986) *J. Mol. Biol.* 189:113-130). Cultures are grown in LB medium containing ampicillin (100 mg/L) at 25°C. At an optical density at 600 nm of approximately 1, IPTG (isopropylthio-β-galactoside, the inducer) can be added to a final concentration of 0.4 mM and incubation can be continued for 3 h at 25°. Cells are then harvested by centrifugation and re-suspended in 50 µL of 50 mM Tris-HCl at pH 8.0 containing 0.1 mM DTT and 0.2 mM phenyl methylsulfonyl fluoride. A small amount of 1 mm glass beads can be added and the mixture sonicated 3 times for about 5 seconds each time with a microprobe sonicator. The mixture is centrifuged and the protein concentration of the supernatant determined. One µg of protein from the soluble fraction of the culture can be separated by SDS-polyacrylamide gel electrophoresis. Gels can be observed for protein bands migrating at the expected molecular weight.

EXAMPLE 7

Assaying Myb-related Transcription Factor Activity

The polypeptides described herein may be produced using any number of methods known to those skilled in the art. Such methods include, but are not limited to, expression in bacteria as described in Example 6, or expression in eukaryotic cell culture, *in planta*, and using viral expression systems in suitably infected organisms or cell lines. The instant -4B. In an alternate embodiment, a thioredoxin fusion protein may be eluted using dithiothreitol; however, elution may be accomplished using other reagents which interact to displace the thioredoxin from the resin. These reagents include β-mercaptoethanol or other reduced

thiol. The eluted fusion protein may be subjected to further purification by polypeptides
may be expressed either as mature forms of the proteins as observed *in vivo* or as fusion
proteins by covalent attachment to a variety of enzymes, proteins or affinity tags. Common
fusion protein partners include glutathione S-transferase ("GST"), thioredoxin ("Trx"),
5 maltose binding protein, and C- and/or N-terminal hexahistidine polypeptide ("His)₆"). The
fusion proteins may be engineered with a protease recognition site at the fusion point so that
fusion partners can be separated by protease digestion to yield intact mature enzyme.
Examples of such proteases include thrombin, enterokinase and factor Xa. However, any
protease can be used which specifically cleaves the peptide connecting the fusion protein and
10 the enzyme.

Purification of the instant polypeptides, if desired, may utilize any number of
separation technologies familiar to those skilled in the art of protein purification. Examples
of such methods include, but are not limited to, homogenization, filtration, centrifugation,
heat denaturation, ammonium sulfate precipitation, desalting, pH precipitation, ion exchange
15 chromatography, hydrophobic interaction chromatography and affinity chromatography,
wherein the affinity ligand represents a substrate, substrate analog or inhibitor. When the
instant polypeptides are expressed as fusion proteins, the purification protocol may include
the use of an affinity resin which is specific for the fusion protein tag attached to the
expressed enzyme or an affinity resin containing ligands which are specific for the enzyme.
20 For example, the instant polypeptides may be expressed as a fusion protein coupled to the
C-terminus of thioredoxin. In addition, a (His)₆ peptide may be engineered into the
N-terminus of the fused thioredoxin moiety to afford additional opportunities for affinity
purification. Other suitable affinity resins could be synthesized by linking the appropriate
ligands to any suitable resin such as Sepharose traditional means as stated above, if desired.
25 Proteolytic cleavage of the thioredoxin fusion protein and the enzyme may be accomplished
after the fusion protein is purified or while the protein is still bound to the ThioBond™
affinity resin or other resin.

Crude, partially purified or purified enzyme, either alone or as a fusion protein, may be
utilized in assays to verify over- or underexpression of functional Myb-related transcription
30 factor protein in transgenic plants and transformed bacterial cells. Assays may be conducted
under well known experimental conditions which permit optimal enzymatic activity. For
example, assays for Myb-related transcription factors are presented by Moyano et al. (1996)
Plant Cell 8:1519-1532.

Various modifications of the invention in addition to those shown and described herein
35 will be apparent to those skilled in the art from the foregoing description. Such
modifications are also intended to fall within the scope of the appended claims.

The disclosure of each reference set forth above is incorporated herein by reference in
its entirety.